

The Star Formation Rate of the Universe at $z \approx 6$ from the Hubble Ultra Deep Field

Andrew J. Bunker^{1,2}, Elizabeth R. Stanway², Richard S. Ellis³,
Richard G. McMahon²

¹ *School of Physics, University of Exeter, Stocker Road, Exeter, EX4 4QL, U.K.*

email: bunker@astro.ex.ac.uk

² *Institute of Astrophysics, University of Cambridge, Madingley Road, Cambridge, CB3 0HA, U.K.*

³ *California Institute of Technology, Mail Stop 169-327, Pasadena, CA 91109, U.S.A.*

Submitted to MNRAS

ABSTRACT

We determine the abundance of i' -band drop-outs in the recently-released *HST*/ACS Hubble Ultra Deep Field (UDF). Since the majority of these sources are likely to be $z \approx 6$ galaxies whose flux decrement between the F775W i' -band and F850LP z' -band arises from Lyman-alpha absorption, the number of detected candidates provides a valuable upper limit to the unextincted star formation rate at this redshift. We demonstrate that the increased depth of UDF enables us to reach an 8σ limiting magnitude of $z'_{AB} = 28.5$ (equivalent to $1.5 h_{70}^{-2} M_{\odot} \text{yr}^{-1}$ at $z = 6.1$, or $0.1 L_{UV}^*$ for the $z \approx 3$ U -drop population), permitting us to address earlier ambiguities arising from the unobserved form of the luminosity function. We identify 54 galaxies (and only one star) at $z'_{AB} < 28.5$ with $(i' - z')_{AB} > 1.3$ over the deepest 11 arcmin^2 portion of the UDF field. The characteristic luminosity (L^*) is consistent with values observed at $z \approx 3$. The faint end slope (α) is less well constrained, but is consistent with only modest evolution. The main change appears to be in the number density (Φ^*). Specifically, and regardless of possible contamination from cool stars and lower redshift sources, the UDF data support our previous result that the star formation rate at $z \approx 6$ was approximately $\times 6$ less than at $z \approx 3$ (Stanway, Bunker & McMahon 2003). This declining comoving star formation rate ($0.005 h_{70} M_{\odot} \text{yr}^{-1} \text{Mpc}^{-3}$ at $z \approx 6$) poses an interesting challenge for models which suggest that the bulk of star forming galaxies that reionized the universe lie at redshifts just beyond $z \simeq 6$.

Key words: galaxies: evolution – galaxies: formation – galaxies: starburst – galaxies: individual: SBM03#1 – galaxies: high redshift – ultraviolet: galaxies

1 INTRODUCTION

There has been considerable progress over the past decade in locating galaxies and QSOs at high redshifts. These sources have enabled us to probe the Universe at early epochs where its physical characteristics are fundamentally different from those at the present epoch. Observations of the most distant $z > 6.2$ QSOs (Becker et al. 2001, Fan et al. 2002) show near-complete absorption at wavelengths shortward of Lyman- α (Gunn & Peterson 1965), suggesting an optical depth in this line that implies a neutral hydrogen fraction which is increasing rapidly with redshift at this epoch. Temperature-polarization cross-correlations in the cosmic microwave background from *WMAP* indicate that the Universe was completely neutral at redshifts of $z > 10$ (Kogut et al. 2003).

Although there is a growing consensus that cosmic

reionization occurred in the redshift interval $6 < z < 15$, a second key question is the nature of the sources responsible for this landmark event. Optical and X-ray studies to $z \simeq 6$ suggest the abundance of active galactic nuclei (AGN) at early epochs is insufficient when account is taken of the relevant unresolved backgrounds (Barger et al. 2003). A more promising source is star-forming galaxies whose early ancestors may be small and numerous. Along with the escape fraction for the ionizing photons from the massive and short-lived OB stars in such sources, a major observational quest in this respect is a determination of the global star formation rate at early epochs.

In previous papers, our group has extended the *Lyman-break* technique (Steidel, Pettini & Hamilton 1995; Steidel et al. 1996) to address this question. Using the Advanced Camera for Surveys (ACS, Ford et al. 2002) on the Hubble Space Telescope (*HST*) with the sharp-sided SDSS F775W

arXiv:astro-ph/0403223 v2 15 Apr 2004

(i') and F850LP (z') filters, we located “ i -drop” candidates with $z'_{AB} < 25.6$ at $z \simeq 6$ for further study. In a series of papers, we have shown that this selection technique can effectively locate $z > 5.7$ galaxies using ACS images from the *HST* Treasury “Great Observatory Origins Deep Survey” (GOODS; Giavalisco & Dickinson 2002). On the basis of GOODS-South photometric catalogues published by Stanway, Bunker & McMahon (2003, hereafter Paper I), spectroscopic follow-up using Keck/DEIMOS and Gemini/GMOS field demonstrated our ability to find high redshift galaxies (Bunker et al. 2003, hereafter Paper II; Stanway et al. 2004a). To address potential cosmic variance issues, we performed a similar analysis in the GOODS-North field, which yielded a consistent estimate of the surface density of $z \simeq 6$ star forming sources (Stanway et al. 2004b, hereafter Paper III).

Although our initial study (Papers I-III) revealed the importance of ascertaining the difficult spectroscopic verifications, and highlighted the problem of contamination from Galactic stars, we nonetheless determined that the abundance of confirmed star forming galaxies at $z \simeq 6$ must be less than that expected on the basis of no evolution from the well-studied $z \sim 3 - 4$ Lyman break population (Steidel et al. 1999). Working at the robustly-detected bright end of the luminosity function, in Paper I we showed that the comoving star formation density in galaxies with $z'_{AB} < 25.6$ is $\approx 6 \times$ less at $z \simeq 6$ than at $z \approx 3$. Our $z'_{AB} < 25.6$ flux limit corresponds to $> 15 h_{70}^{-2} M_{\odot} \text{yr}^{-1}$ at $z = 5.9$, equivalent to L_{UV}^* at $z \approx 3$. Other groups have claimed less dramatic evolution or even no evolution in the volume-averaged star formation rate, based on the same fields (Giavalisco et al. 2004; Dickinson et al. 2004) and similar *HST*/ACS data sets (Bouwens et al. 2003; Yan, Windhorst & Cohen 2003), but working closer to the detection limit of the images and introducing large completeness corrections for the faint source counts. The major uncertainty in converting the abundance of our spectroscopically-confirmed sample in the GOODS fields into a $z \simeq 6$ comoving star formation rate is the form of the luminosity function for faint, unobserved, sources. As discussed in Paper III, if the faint end of the luminosity function at $z \simeq 6$ was steeper than that at lower redshift, or if L^* was significantly fainter, a non-evolving star formation history could perhaps still be retrieved.

The public availability of the Hubble Ultra Deep Field (UDF; Beckwith, Somerville & Stiavelli 2003) enables us to address this outstanding uncertainty. By pushing the counts and the inferred luminosity function of i' -band drop-outs at $z \approx 6$ to a limiting lower luminosity equivalent to one well below L_3^* for the $z \approx 3$ population, it is possible to refine the integrated star formation rate at $z \approx 6$. In this paper we set out to undertake the first photometric analysis of i' -drops in the UDF. Our primary goal is to understand the abundance of fainter objects with characteristics equivalent to those of $z \simeq 6$ sources and address uncertainties in the global star formation rate at this redshift.

The structure of the paper is as follows. In Section 2 we describe the imaging data, the construction of our catalogues and our i' -drop selection. In Section 3 we discuss the luminosity function of star-forming sources, likely contamination on the basis of earlier spectroscopic work, and estimate the density of star formation at $z \simeq 6$. Our conclusions are presented in Section 4. Throughout we adopt the

standard “concordance” cosmology of $\Omega_M = 0.3$, $\Omega_{\Lambda} = 0.7$, and use $h_{70} = H_0/70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. All magnitudes are on the AB system (Oke & Gunn 1983).

2 HST IMAGING: OBSERVATIONS AND I -DROP SELECTION

2.1 *HST*/ACS Observations

The Hubble Ultra Deep Field (UDF) is a public *HST* survey made possible by Cycle 12 STScI Director’s Discretionary Time programme GO/DD-9978 executed over September 2003 – January 2004. For the present program, the *HST* has imaged a single ACS Wide Field Camera (WFC) tile (11.5 arcmin^2) for 400 orbits in 4 broad-band filters (F435W B -band for 56 orbits; F606W V -band for 56 orbits; F775W i' -band for 144 orbits; F850LP z' -band for 144 orbits). The UDF field lies within the Chandra Deep Field South (CDF-S) with coordinates RA= $03^{\text{h}}32^{\text{m}}39^{\text{s}}.0$, Decl.= $-27^{\circ}47'29''.1$ (J2000). As the UDF represents the deepest set of images yet taken, significantly deeper than the I -band exposures of the Hubble Deep Fields (Williams et al. 1996; 1998), and adds the longer-wavelength z' -band, it is uniquely suited to the goals of our program.

The WFC on ACS has a field of $202'' \times 202''$, and a pixel scale of $0.05''$. The UDF lies within the survey area of GOODS-South area (Giavalisco et al. 2004), surveyed using ACS with the same filters to shallower depth (3,2.5,2.5 & 5 orbits in the B , V , i' & z' bands). The UDF was observed at two main orientations differing by 90 degrees, and within each of these data was taken in 2 blocks rotated by 4 deg (orientations of 310,314,40 & 44 deg). A 4-point dither box spanning 0.3 arcsec was used, with half-pixel centres to improve the sampling. During each “visit”, there were 3 larger 3 arcsec dithers to span the WFC inter-chip gap.

For our analysis we use the reduced UDF data v1.0 made public by STScI on 09 March 2004. The pipeline reduction involved bias/dark current subtraction, flat-fielding, and the combination of background-subtracted frames rejecting cosmic ray strikes and chip defects. The resulting reduced images had been “drizzled” (Fruchter & Hook 2002) using the “MultiDrizzle” software (Koekemoer et al. 2004) on to a finer pixel scale of $0.03''$, to correct for geometric distortion and to improve the sampling of the point spread function (PSF). The UDF data has been placed on the same astrometric system as the GOODSv1.0 images of the UDF¹. The photometric zeropoints adopted were those provided by STScI for the UDF v1.0 data release: 25.673, 26.486, 25.654 & 24.862 for the B , V , i' & z' filters, where $\text{mag}_{AB} = \text{zeropoint} - 2.5 \log_{10}(\text{counts/s})$. We have corrected for the small amount of foreground Galactic extinction toward the CDFS using the *COBE*/DIRBE & *IRAS*/ISSA dust maps of Schlegel, Finkbeiner & Davis (1998). The optical reddening is $E(B - V) = 0.008$, equivalent to extinctions of $A_{F775} = 0.017$ & $A_{F850LP} = 0.012$.

¹ Available from <ftp://archive.stsci.edu/pub/hlsp/goods/v1>

2.2 Construction of Catalogues

Candidate selection for all objects in the field was performed using version 2.3.2 of the SExtractor photometry package (Bertin & Arnouts 1996). As we are searching specifically for objects which are only securely detected in z' , with minimal flux in the i' -band, fixed circular apertures $0''.5$ in diameter were trained in the z' -image and the identified apertures used to measure the flux at the same spatial location in the i' -band image by running SExtractor in two-image mode. For object identification, we adopted a limit of at least 5 contiguous pixels above a threshold of 2σ per pixel (0.0005 counts/pixel/s) on the data drizzled to a scale of $0''.03$ pixel $^{-1}$. This cut enabled us to detect all significant sources and a number of spurious detections close to the noise limit. As high redshift galaxies in the rest-UV are known to be compact (e.g., Ferguson et al. 2004, Bremer et al. 2004), we corrected the aperture magnitudes to approximate total magnitudes through a fixed aperture correction, determined from bright compact sources: -0.11 mag in i' -band and -0.14 mag in z' -band, the larger latter correction arising from the more extended PSF wings of the z' -band.

The measured noise in the drizzled images underestimates the true noise as adjacent pixels are correlated. To assess the true detection limit and noise properties, we examined the raw ACS/WFC images from the *HST* archive and measured the noise in statistically-independent pixels. For the 144-orbit z' -band, we determine that the 8σ detection limit is $z'_{AB} = 28.5$ for our $0''.5$ -diameter aperture. This is consistent with the noise decreasing as $\sqrt{\text{time}}$ from the 5-orbit GOODSv1.0 to the 144-orbit UDF z' -band. We adopt this high $S/N = 8$ cut as our conservative sample limit. We trimmed the outermost edges where fewer frames overlapped in order to exploit the deepest UDF region, corresponding to a survey area of 11 arcmin 2 . From the output of SExtractor ($\simeq 60,000$ objects in our z' -band selected catalogue, many spurious close to the detection limit) we created a sub-catalogue of all real objects brighter than $z'_{AB} < 28.5$ mag (8σ in a $0''.5$ -diameter aperture), of which 63 appear to be promising i' -band dropouts (see Section 2.3) with $(i' - z')_{AB} > 1.3$.

To quantify possible incompleteness in this catalogue, we adopted two approaches. First we examined the recovery rate of artificial galaxies created with a range of total magnitudes and sizes. We used de Vaucouleurs $r^{1/4}$ and exponential disk profiles, convolved with the ACS/WFC PSF derived from unsaturated stars in the UDF images. Secondly we created fainter realisations of the brightest i' -dropout in the UDF confirmed to be at high redshift (SBM03#1 with $z'_{AB} = 25.4$, confirmed spectroscopically to be at $z = 5.83$ by Stanway et al. 2004b; Dickinson et al. 2004). By excising a small region around this i' -dropout, scaling the sub-image to a fainter magnitude, and adding it back into the UDF data at random locations, we assessed the recoverability as a function of brightness. For such objects we recover 98% of the simulated sources to $z'_{AB} = 28.5$, the remainder being mainly lost via source confusion through overlapping objects. From these analyses, we determine that, for unresolved sources ($r_h = 0''.05$), we are complete at our 8σ limit of $z'_{AB} = 28.5$, and are 97% complete at this magnitude for $r_h = 0''.2$ (Figure 1). For objects with larger half-light radii we will underestimate the z' -band flux due to our $0''.5$ -




Figure 1. The completeness (normalized to unity) for artificial galaxies added to the UDF z' -band image, as a function of total magnitude and half-light radius; we re-ran SExtractor on this image to assess the fraction of artificial galaxies recovered. The completeness is $> 97\%$ for $R_h < 0''.2$ and $z'_{AB} < 28.5$.

diameter photometric aperture. However, this effect is small for our sample of compact sources (Table 1 lists both the $0''.5$ -diameter magnitudes with an aperture correction which we adopt, and the SExtractor “AUTO” estimate of the total magnitude: these are broadly consistent).

At the relatively bright cut of $z'_{AB} < 25.6$ used in Paper I from the GOODSv0.5 individual epochs, the UDF data is 98% complete for sources as extended as $r_h = 0.5$ arcsec. Interestingly, we detect no extended (low surface brightness) i' -drops to this magnitude limit in addition to SBM03#1 (Papers I,III) in the deeper UDF data. This supports our assertion (Paper I) that the i' -drop population is predominantly compact and there cannot be a large completeness correction arising from extended objects (c.f. Lanzetta et al. 2002). The ACS imaging is of course picking out HII star forming regions, and these UV-bright knots of star formation are typically < 1 kpc even within large galaxies at low redshift.

2.3 Redshift Discrimination

In order to select $z > 6$ galaxies, we use the Lyman break technique pioneered at $z \sim 3$ using ground-based telescopes by Steidel and co-workers and using *HST* by Madau et al. (1996). At $z \sim 3 - 4$ the technique involves the use of three filters: one below the Lyman limit ($\lambda_{\text{rest}} = 912 \text{ \AA}$), one in the Lyman forest region and a third longward of the Lyman- α line ($\lambda_{\text{rest}} = 1216 \text{ \AA}$). At $z \simeq 6$, we can efficiently use only two filters, since the integrated optical depth of the Lyman- α forest is $\gg 1$ (see Figure 2) rendering the shortest-wavelength filter below the Lyman limit redundant. The key issue is to work at a sufficiently-high signal-to-noise ratio that i' -band drop-outs can be safely identified through de-

tection in a single redder band (i.e., SDSS- z'). This approach has been demonstrated to be effective by the SDSS collaboration in the detection of $z \sim 6$ quasars using the i' - and z' -bands alone (Fan et al. 2001). The sharp sides of the SDSS filters assist in the clean selection using the photometric redshift technique. In Figures 3 & 4 we illustrate how a colour cut of $(i' - z')_{AB} > 1.5$ (used in Papers I-III) can be effective in selecting sources with $z > 5.7$. Here we relax this cut to $(i' - z')_{AB} > 1.3$ to recover most galaxies at redshifts $z > 5.6$, but at the expense of potentially larger contamination by $z \approx 1 - 2$ ellipticals. Near-infrared colours from the NICMOS imaging of the UDF should identify these Extremely Red Objects (EROs), and we consider this in a companion paper (Stanway, McMahon & Bunker 2004c).

Six of the 63 candidate i' -dropouts in our $z'_{AB} < 28.5$ UDF catalogue were identified visually as different regions of the same extended source, and where these were within our aperture diameter of $0''.5$ the duplicates were eliminated from the final selection. One spurious i' -drop arose from the diffraction spike of a bright star due to the more extended PSF in the z' -band compared with that in the i' -band. Only one of the i' -dropouts is unresolved (Figure 5). This is the brightest at $z'_{AB} = 25.3$ (#11337 in Table 1), detected in the V -band image and removed from our catalogue of potential $z \approx 6$ objects as a probable star. At the edge of the UDF frame (and outside the central 11 arcmin^2 region of lowest noise where we do main our analysis) there is a second unresolved i' -drop with $z'_{AB} = 25.2$, first identified in Paper I (SBM03#5), where we argued that the near-IR colours are likely to be stellar. It is interesting that the level of stellar contamination in the UDF i' -drops is only 2%, compared with about one in three at the bright end ($z'_{AB} < 25.6$, Papers I & III). This may be because we are seeing through the Galactic disk at these faint magnitudes to a regime where there are no stars at these faint limiting magnitudes.

From our original list of 63 i' -drops, 6 duplications were removed, along with one diffraction spike artifact. The remaining objects satisfying our $(i' - z')_{AB} > 1.3$ & $z'_{AB} < 28.5$ selection criteria are detailed in Table 1, of which 54 are good candidate $z \approx 6$ galaxies, along with the probable star #11337, and another objected (#46574) detected in V -band. The surface density of i' -drops as a function of limiting magnitude is shown in Figure 7.

3 THE LUMINOSITY FUNCTION OF STAR FORMING GALAXIES AT $Z \approx 6$

3.1 Estimate of Star Formation Rate from the Rest-UV

We will base our measurement of the star formation rate for each candidate on the rest-frame UV continuum, redshifted into the z' -band at $z \approx 6$ and measured from the counts in a $0''.5$ -diameter aperture (with an aperture correction to total magnitudes, Section 2.2). In the absence of dust obscuration, the relation between the flux density in the rest-UV around $\approx 1500 \text{ \AA}$ and the star formation rate (SFR in $M_{\odot} \text{ yr}^{-1}$) is given by $L_{UV} = 8 \times 10^{27} \text{ SFR ergs s}^{-1} \text{ Hz}^{-1}$ from Madau, Pozzetti & Dickinson (1998) for a Salpeter (1955) stellar initial mass function (IMF) with $0.1 M_{\odot} < M^* < 125 M_{\odot}$. This is comparable to the relation derived from the models of Leitherer & Heckman (1995) and Kennicutt (1998).




Figure 2. The ACS- i' and $-z'$ bandpasses overplotted on the spectrum of a generic $z = 6$ galaxy (solid line), illustrating the utility of our two-filter technique for locating $z \approx 6$ sources.

However, if a Scalo (1986) IMF is used, the inferred star formation rates will be a factor of ≈ 2.5 higher for a similar mass range.

Recognising the limitations of our earlier studies (Papers I-III) which by necessity focussed on the brighter i' -drops, we now attempt to recover the $z \approx 6$ rest-frame UV luminosity function from the observed number counts of i' -drops to faint magnitudes in the UDF. Although our colour cut selects galaxies with redshifts in the range $5.6 < z < 7.0$, an increasing fraction of the z' -band flux is attenuated by the redshifted Lyman- α forest. At higher redshifts we probe increasingly shortward of $\lambda_{\text{rest}} = 1500 \text{ \AA}$ (where the luminosity function is calculated) so the k -corrections become significant beyond $z \approx 6.5$.

Figure 6 demonstrates this bias and shows the limiting star formation rate as a function of redshift calculated by accounting for the filter transmissions and the blanketing effect of the intervening Lyman- α forest. By introducing the small k -correction to $\lambda_{\text{rest}} = 1500 \text{ \AA}$ from the observed rest-wavelengths longward of Lyman- α redshifted into the z' -band we can correct for this effect. We considered a spectral slope of $\beta = -2.0$ (where $f_{\lambda} \propto \lambda^{\beta}$) appropriate for an unobscured starburst (flat in f_{ν}), and also a redder slope of $\beta = -1.1$ which appropriate for mean reddening of the $z \approx 3$ Lyman break galaxies given by Meurer et al. (1997). A more recent determination for this population by Adelberger & Steidel (2000) gives $\beta = -1.5$, in the middle of the range. At our 8σ limiting magnitude of $z'_{AB} = 28.5$, we deduce we can detect unobscured star formation rates as low as $1.0 [1.1] h_{70}^{-2} M_{\odot} \text{ yr}^{-1}$ at $5.6 < z < 5.8$ and $1.5 [1.7] h_{70}^{-2} M_{\odot} \text{ yr}^{-1}$ at $z < 6.1$ for spectral slope $\beta = -2.0 [-1.1]$ (Figure 6).

Recognising that contamination by interlopers will only reduce the value, we now compare the comoving star formation rate deduced for $z \approx 6$ galaxies based on our can-

Table 1. i' -band dropouts in the UDF. The two stars are above the line – all others are spatially resolved. Our ID and the corresponding match from the UDF catalogues released by STScI are listed in columns one and two. Where two i' -drops are close, and lie within our $0''.5$ -diameter aperture, they are indicated and the flux only counted once in the star formation total – those IDs and star formation rates in parentheses are not counted. The star formation rates assume the i' -drops lie at $z = 6.0$, the expected median redshift of our sample.

Our ID	STScI ID	RA & Declination (J2000)			z'_{AB} ($0''.5$ -diameter aperture)	i'_{AB}	$(i' - z')_{AB}$ $0''.5$ -aper	R_h arcsec	z'_{AB} (total)	$\text{SFR}_{\text{UV}}^{z=6}$ $h_{70}^{-2} M_{\odot} \text{yr}^{-1}$
(2140)*	—	03 32 38.80	−27 49 53.6		25.22 ± 0.02	27.91 ± 0.04	2.69 ± 0.05	0.06	25.17 ± 0.02	(star)
(11337)	443	03 32 38.02	−27 49 08.4		25.29 ± 0.02	26.79 ± 0.04	1.50 ± 0.05	0.05	25.43 ± 0.02	(star)
20104 ¹	2225	03 32 40.01	−27 48 15.0		25.35 ± 0.02	26.99 ± 0.03	1.64 ± 0.04	0.08	25.29 ± 0.02	$19.5[z = 5.83]$
42929 ²	8033	03 32 36.46	−27 46 41.4		26.56 ± 0.03	29.05 ± 0.14	2.49 ± 0.15	0.14	26.55 ± 0.04	6.75
41628	8961	03 32 34.09	−27 46 47.2		26.65 ± 0.04	28.81 ± 0.12	2.15 ± 0.12	0.09	26.70 ± 0.04	6.18
(46574) ³	7730	03 32 38.28	−27 46 17.2		26.71 ± 0.04	29.38 ± 0.18	2.67 ± 0.18	0.09	26.74 ± 0.04	(5.87)
24019	3398	03 32 32.61	−27 47 54.0		26.80 ± 0.04	28.22 ± 0.08	1.42 ± 0.09	0.18	26.73 ± 0.04	5.42
52880	9857	03 32 39.07	−27 45 38.8		27.00 ± 0.05	28.47 ± 0.09	1.47 ± 0.10	0.09	27.10 ± 0.05	4.50
23516	3325	03 32 34.55	−27 47 56.0		27.04 ± 0.05	28.57 ± 0.10	1.53 ± 0.11	0.11	27.05 ± 0.05	4.35
10188	322	03 32 41.18	−27 49 14.8		27.10 ± 0.05	29.15 ± 0.16	2.04 ± 0.16	0.20	27.06 ± 0.05	4.10
21422	2690	03 32 33.78	−27 48 07.6		27.23 ± 0.05	28.99 ± 0.14	1.76 ± 0.15	0.10	27.37 ± 0.05	3.64
25578 ^D	—	03 32 47.85	−27 47 46.4		27.30 ± 0.06	29.96 ± 0.31	2.66 ± 0.31	0.18	27.28 ± 0.06	3.41
25941	4050	03 32 33.43	−27 47 44.9		27.32 ± 0.06	29.30 ± 0.18	1.99 ± 0.19	0.11	27.38 ± 0.06	3.35
26091 ^D	4110	03 32 41.57	−27 47 44.2		27.38 ± 0.06	29.74 ± 0.25	2.35 ± 0.26	0.14	27.21 ± 0.07	3.16
24458	3630	03 32 38.28	−27 47 51.3		27.51 ± 0.07	29.11 ± 0.15	1.60 ± 0.17	0.18	27.67 ± 0.08	2.80
21262	2624	03 32 31.30	−27 48 08.3		27.52 ± 0.07	28.96 ± 0.13	1.44 ± 0.15	0.20	27.49 ± 0.08	2.78
13494	30591	03 32 37.28	−27 48 54.6		27.56 ± 0.07	30.62 ± 0.55	3.06 ± 0.55	0.12	27.48 ± 0.08	2.69
24228	3450	03 32 34.28	−27 47 52.3		27.63 ± 0.07	29.10 ± 0.15	1.47 ± 0.17	0.17	27.39 ± 0.08	2.52
16258	1400	03 32 36.45	−27 48 34.3		27.64 ± 0.07	29.07 ± 0.15	1.42 ± 0.16	0.18	27.25 ± 0.07	2.49
42414	9202	03 32 33.21	−27 46 43.3		27.65 ± 0.07	29.10 ± 0.15	1.45 ± 0.17	0.16	27.54 ± 0.08	2.46
27173 ⁵	4377	03 32 29.46	−27 47 40.4		27.73 ± 0.08	29.87 ± 0.28	2.13 ± 0.29	0.13	27.74 ± 0.09	2.28
49117 ^D	—	03 32 38.96	−27 46 00.5		27.74 ± 0.08	29.77 ± 0.26	2.03 ± 0.27	0.17	27.36 ± 0.07	2.28
49701	36749	03 32 36.97	−27 45 57.6		27.78 ± 0.08	30.79 ± 0.64	3.02 ± 0.64	0.19	27.90 ± 0.09	2.20
24123	—	03 32 34.29	−27 47 52.8		27.82 ± 0.08	29.89 ± 0.29	2.07 ± 0.30	0.15	27.65 ± 0.09	2.11
27270	33003	03 32 35.06	−27 47 40.2		27.83 ± 0.08	30.69 ± 0.58	2.87 ± 0.59	0.11	27.99 ± 0.09	2.10
23972	3503	03 32 34.30	−27 47 53.6		27.84 ± 0.09	29.38 ± 0.19	1.54 ± 0.21	0.17	27.77 ± 0.10	2.07
14751	1086	03 32 40.91	−27 48 44.7		27.87 ± 0.09	29.27 ± 0.17	1.40 ± 0.19	0.09	27.92 ± 0.09	2.02
44154	35945	03 32 37.46	−27 46 32.8		27.87 ± 0.09	$> 30.4 (3\sigma)$	$> 2.5 (3\sigma)$	0.16	27.87 ± 0.10	2.01
35084	34321	03 32 44.70	−27 47 11.6		27.92 ± 0.09	29.86 ± 0.28	1.94 ± 0.30	0.14	27.90 ± 0.09	1.93
42205	8904	03 32 33.55	−27 46 44.1		27.93 ± 0.09	29.51 ± 0.21	1.57 ± 0.23	0.11	27.91 ± 0.09	1.90
46503	7814	03 32 38.55	−27 46 17.5		27.94 ± 0.09	29.43 ± 0.20	1.50 ± 0.22	0.12	28.07 ± 0.09	1.89
19953	2225	03 32 40.04	−27 48 14.6		27.97 ± 0.09	29.50 ± 0.21	1.54 ± 0.23	0.17	27.68 ± 0.10	1.85
52086	36786	03 32 39.45	−27 45 43.4		27.97 ± 0.09	30.83 ± 0.66	2.86 ± 0.66	0.11	28.04 ± 0.10	1.84
44194	35945	03 32 37.48	−27 46 32.5		28.01 ± 0.10	30.61 ± 0.54	2.60 ± 0.55	0.18	27.46 ± 0.09	1.77
21111 ^D	2631	03 32 42.60	−27 48 08.9		28.02 ± 0.10	29.69 ± 0.24	1.67 ± 0.26	0.15	28.08 ± 0.10	1.76
46223 ⁴	35506	03 32 39.87	−27 46 19.1		28.03 ± 0.10	32.18 ± 2.23	4.15 ± 2.23	0.14	28.10 ± 0.11	1.74
22138	32007	03 32 42.80	−27 48 03.2		28.03 ± 0.10	$> 30.4 (3\sigma)$	$> 2.3 (3\sigma)$	0.14	28.14 ± 0.10	1.73
(46234) ⁴	—	03 32 39.86	−27 46 19.1		28.05 ± 0.10	30.61 ± 0.54	2.56 ± 0.55	0.12	28.30 ± 0.12	(1.70)
14210	978	03 32 35.82	−27 48 48.9		28.08 ± 0.10	29.51 ± 0.21	1.43 ± 0.24	0.10	28.16 ± 0.11	1.66
45467	35596	03 32 43.02	−27 46 23.7		28.08 ± 0.10	$> 30.4 (3\sigma)$	$> 2.3 (3\sigma)$	0.11	28.25 ± 0.10	1.66
12988 ^D	30534	03 32 38.49	−27 48 57.8		28.11 ± 0.11	30.47 ± 0.48	2.36 ± 0.49	0.10	28.22 ± 0.11	1.61
30359	33527	03 32 30.14	−27 47 28.4		28.13 ± 0.11	29.58 ± 0.22	1.46 ± 0.25	0.13	28.02 ± 0.11	1.59
11370	482	03 32 40.06	−27 49 07.5		28.13 ± 0.11	30.45 ± 0.47	2.32 ± 0.48	0.06	28.27 ± 0.08	1.59
24733	32521	03 32 36.62	−27 47 50.0		28.15 ± 0.11	30.92 ± 0.71	2.76 ± 0.72	0.13	28.34 ± 0.12	1.55
37612	34715	03 32 32.36	−27 47 02.8		28.18 ± 0.11	29.98 ± 0.31	1.80 ± 0.33	0.13	28.15 ± 0.11	1.52
41918	7829	03 32 44.70	−27 46 45.5		28.18 ± 0.11	29.81 ± 0.27	1.63 ± 0.29	0.08	28.36 ± 0.10	1.52
21530	31874	03 32 35.08	−27 48 06.8		28.21 ± 0.12	30.24 ± 0.39	2.03 ± 0.41	0.12	28.35 ± 0.12	1.47
42806	8033	03 32 36.49	−27 46 41.4		28.21 ± 0.12	30.76 ± 0.62	2.55 ± 0.63	0.11	28.12 ± 0.11	1.47
27032 ⁵	4377	03 32 29.45	−27 47 40.6		28.22 ± 0.12	29.55 ± 0.22	1.34 ± 0.25	0.06	28.70 ± 0.12	1.46
52891	36697	03 32 37.23	−27 45 38.4		28.25 ± 0.12	32.21 ± 2.28	3.96 ± 2.28	0.16	28.34 ± 0.11	1.43
17908	1834	03 32 34.00	−27 48 25.0		28.25 ± 0.12	29.66 ± 0.24	1.41 ± 0.27	0.15	28.22 ± 0.13	1.42
(27029) ⁵	4353	03 32 29.44	−27 47 40.7		28.25 ± 0.12	29.98 ± 0.31	1.73 ± 0.33	0.09	28.67 ± 0.14	(1.42)
48989 ^D	36570	03 32 41.43	−27 46 01.2		28.26 ± 0.12	$> 30.4 (3\sigma)$	$> 2.1 (3\sigma)$	0.09	28.45 ± 0.12	1.41
17487	—	03 32 44.14	−27 48 27.1		28.30 ± 0.12	30.10 ± 0.35	1.81 ± 0.37	0.07	28.51 ± 0.11	1.36
18001	31309	03 32 34.14	−27 48 24.4		28.40 ± 0.13	30.46 ± 0.48	2.06 ± 0.49	0.14	28.59 ± 0.14	1.23
35271	6325	03 32 38.79	−27 47 10.9		28.44 ± 0.14	29.77 ± 0.26	1.33 ± 0.30	0.10	28.60 ± 0.13	1.19
22832	—	03 32 39.40	−27 47 59.4		28.50 ± 0.15	30.46 ± 0.47	1.96 ± 0.50	0.14	28.60 ± 0.13	1.13

^D double. * star SBM03# 5 (Paper I), outside central UDF. ¹ SBM03#1 (Paper I); SiD002 (Dickinson et al. 2004). ² SiD025 (Dickinson et al. 2004). ³ 46574 has a close neighbour visible in the v -band (i.e. low redshift.) ⁴ 46234 is close to 46223. ⁵ 27029 is close to 27032.




Figure 3. Model colour-redshift tracks for galaxies with non-evolving stellar populations (from Coleman, Wu & Weedman 1980 template spectra). The contaminating ‘hump’ in the $(i' - z')$ colour at $z \approx 1 - 2$ arises when the Balmer break and/or the 4000 Å break redshifts beyond the i' -filter.

didate i' -dropout source counts with predictions based on a range of rest-frame UV luminosity functions. For convenience we assume that there is no evolution over the sampled redshift range, $5.6 < z < 6.5$, spanned by the UDF data (equivalent to a range between $0.8 - 1.0 h_{70}^{-1}$ Gyr after the Big Bang). We take as a starting point the luminosity function for the well-studied Lyman-break U -dropout population, reported in Steidel et al. (1999), which has a characteristic rest-UV luminosity $m_R^* = 24.48$ (equivalent to $M_3^*(1500 \text{ \AA}) = -21.1$ mag or $L_3^* = 15 h_{70}^{-2} M_\odot \text{ yr}^{-1}$ for our cosmology). The faint end slope of the Schechter function at $z \approx 3$ is relatively steep ($\alpha = -1.6$) compared with $\alpha = -1.0$ to -1.3 for lower-redshift galaxy samples (e.g., Lilly et al. 1995; Efstathiou et al. 1988; Blanton et al. 2003 – see Gabasch et al. 2004 for recent determinations at 1500 Å). The characteristic comoving number density at $z \approx 3$ is $\Phi_3^* = 0.00138 h_{70}^3 \text{ Mpc}^{-3}$ in our cosmology.

We constructed a grid of models based upon the $z \approx 3$ luminosity function, varying α between -1.1 and -1.9 , and L^* between $0.3 L_3^*$ and $2 L_3^*$. Leaving the normalization of the luminosity function, Φ^* , as a free parameter, we model the effect of the break below the Lyman- α emission line due to blanketing by the forest, where the continuum break D_A (Oke & Korycansky 1982) is defined as

$$D_A = \left(1 - \frac{f_\nu(1050 - 1170 \text{ \AA})_{\text{obs}}}{f_\nu(1050 - 1170 \text{ \AA})_{\text{pred}}} \right). \quad (1)$$

We assumed flux decrements of $D_A = 0.9 - 1.0$, consistent with that observed in the $z > 5.8$ SDSS QSOs (Fan et al. 2001).

We find that altering the spectral slope β changes the predicted number of i' -dropouts by only $\approx 10\%$, although lowering D_A reduces the completeness in the lowest redshift




Figure 4. Colour-magnitude diagram for the UDF data with the limit $z'_{AB} < 28.5$ and $(i' - z')_{AB} = 1.3$ colour cut shown (dashed lines). As discussed in the text, such a catalogue could be contaminated by cool stars, EROs and wrongly identified extended objects and diffraction spikes but nonetheless provides a secure upper limit to the abundance of $z \approx 6$ star forming galaxies. Circles and arrows (lower limits) indicate our i' -drop candidate $z \approx 6$ galaxies. The solid circle is the spectroscopically-confirmed galaxy SBM03#1 (Stanway et al. 2004b; Dickinson et al. 2004), and the asterisk is the only unresolved i' -drop in our UDF sample, the probable star #11337.

bin $5.6 < z < 5.8$ for a $(i' - z')_{AB} > 1.5$ colour cut. A $(i' - z')_{AB} > 1.3$ cut improves the selection somewhat but at the risk of higher contamination from red objects at $z \approx 1 - 2$: we consider this in Stanway, McMahon & Bunker (2004c).

We minimize χ^2 for our grid of model luminosity functions: our best fit (Figure 7) is compatible with no evolution of L^* from $z \approx 3$, but a large decline in the comoving space density, Φ^* (by about a factor 6 relative to $z \approx 3$). The faint end slope is less well constrained, although no evolution is compatible with the results. At the faintest magnitude bin, there are slightly higher counts, perhaps indicating a slightly steeper α if the results at the faintest magnitudes are to be trusted (Figure 8).

For the simple no-evolution model (using the same luminosity function at $z = 6$ as at $z = 3$), 170 galaxies satisfying our $i'_{AB} < 28.5$ & $(i' - z')_{AB} > 1.3$ selection with a total star formation rate of $870 h_{70}^{-2} M_\odot \text{ yr}^{-1}$ would be predicted (with faint-end slope $\alpha = -1.6$, spectral slope $\beta = -2.0$ and Lyman- α forest decrement $D_A = 1.0$). This compares with our observed 54 i' -drops (1/3rd the predicted number), which would have a total star formation rate of $140 h_{70}^{-2} M_\odot \text{ yr}^{-1}$ (1/6th of the no-evolution prediction). A calculation using the effective volume (V_{eff} , see Steidel et al. 1999) as in Paper I yields the same result as the comparison here with a simulated non-evolving population with the $z \approx 3$ parameters. The predicted median redshift of our




Figure 5. The distribution of angular sizes (half-light radius, R_h , in arcseconds) for objects in our z' -band selected catalogue. Our i' -drop candidate $z \approx 6$ are marked as open circles, with the confirmed $z = 5.8$ galaxy SBM03#1 a solid circle. The i' -drops appear to be compact but resolved (the stellar locus at $0''.05$ is clearly visible). The asterisk denotes the only unresolved i' -drop in our UDF sample, the probable star #11337.

i' -drop sample for the no-evolution model is $z = 5.95$, with the luminosity-weighted average $\bar{z} = 6.05$.

Recognizing the very limited area of the UDF and the problems of cosmic variance, it is nonetheless interesting to compare our measured i' -drop number counts with previous determinations from shallower data sets. The surface density derived in Paper I to $z'_{AB} = 25.6$ is consistent with the present data – we detect only one resolved i' -dropout this bright: SBM03#1. Note that the UDF pointing was selected to include this object. No other spatially-resolved i' -dropouts are detected to $z'_{AB} < 26.5$, implying a surface density of $0.1 \pm 0.1 \text{ arcmin}^{-2}$. This is in contrast with the density of 0.4 arcmin^{-2} measured by Giavalisco et al. (2004), and $0.5 \pm 0.2 \text{ arcmin}^{-2}$ from the completeness-corrected estimate of Bouwens et al. (2003)², and the even higher surface density of 2.3 arcmin^{-2} claimed by Yan, Windhorst & Cohen (2003), after correcting for a factor of 4 error in their

² *Note added in proof:* a recent paper by Bouwens et al. (2004), based on number counts of i' -drops in the ACS parallel observations to the NICMOS UDF field, significantly revises their previous estimate of the number density of $z'_{AB} < 28.5$ i' -drops from 0.5 ± 0.2 to $0.2 \pm 0.1 \text{ arcmin}^{-2}$ (4 objects in 21 arcmin^2 , consistent with our UDF work presented here). The conclusion of Bouwens et al. (2003) –that the comoving star formation at $z \approx 6$ is consistent with no evolution from $z \approx 4$ – is revised in Bouwens et al. (2004) to be a factor of 3 decline from $z = 3.8$ to $z \approx 6$. Using the evolution in comoving number density of $(1+z)^{-2.8}$ suggested by Bouwens et al. (2004), this fall in star formation rate at $z = 6$ is consistent with our result of a factor of 6 decline from $z = 3.0$ to $z \approx 6$ from the GOODS data in Stanway, Bunker & McMahon (2003), confirmed in this paper from the deeper UDF data.




Figure 6. Limiting star formation rate as a function of redshift for the UDF catalogue with $z'_{AB} < 28.5$ mag (8σ). Star formation rates are inferred from the rest-frame 1500 \AA flux (Madau, Pozzetti & Dickinson 1998) taking account of k -corrections, filter transmission and blanketing by Lyman- α absorption. The solid line assumes a spectral slope $\beta = -2.0$ (where $f_\lambda \propto \lambda^\beta$) appropriate for an unobscured starburst, and the dotted line has $\beta = -1.1$ (corresponding to mean reddening of $z \approx 3$ Lyman break galaxies given in Meurer et al. 1997). The limit as a fraction of L_3^* ($L^*[1500\text{\AA}]$ at $z \approx 3$, equivalent to $\text{SFR}_{UV}^* = 15 h_{70}^{-2} M_\odot \text{ yr}^{-1}$ from Steidel et al. 1999) is shown on the right axis. Our colour selection should remove most $z < 5.6$ galaxies (solid vertical line), and our average i' -drop redshift for $z' < 28.5$ should be $z \approx 6.0$ (vertical dot-dash line): we are sensitive as faint at $0.1 L_3^*$ at this redshift.

original flux calibration (see Yan & Windhorst 2004). The discrepancies may be due to cosmic variance, or too many spurious sources in the samples of these teams, due to working close to the sensitivity limits.

From Somerville et al. (2004) we estimate that the cosmic variance for the UDF is 40%, assuming the $z = 6$ LBGs are clustered in the same way as the $z = 3$ LBGs and assuming a volume of derived by scaling our UDF area with our wider-area GOODS data (with an effective volume of $1.8 \times 10^5 h_{70}^{-3} \text{ Mpc}^3$ for the 146 arcmin^2 of GOODS-S, Paper I). Indeed, the spatial distribution of our i' -drops on the sky does indicate some clustering (Figure 9), and we had already flagged 6 of our candidates as being “double” sources (Table 1), with another 2 having near neighbours. In the GOODS survey of the CDF-S, Stanway et al. (2004a) have already spectroscopically identified an overdensity at $z = 5.8$.

3.2 Implications for Reionization

The increased depth of the UDF enables us to resolve the uncertainties associated with the unobserved portion of the luminosity function (LF) for $z \approx 6$ sources. Our best-fit LF suggest little or no change in L^* and α over $3 < z < 6$




Figure 7. Cumulative source counts per arcmin² of i' -dropout as a function of z' -band magnitude. The new UDF data (over a smaller area of 11 arcmin² for $z'_{AB} > 27.0$) is compared with $z'_{AB} < 25.6$ single epoch GOODSv0.5 ACS/WFC imaging over 300 arcmin² (Papers I-III) and combined 5 epoch GOODSv1.0 images to $z'_{AB} < 27.0$ (Stanway 2004).




Figure 8. Cumulative source counts per arcmin² of i' -dropout as a function of z' -band magnitude, with various values of the faint end slope (α) assuming $L_6^* = L_3^*$ and $\Phi_6^* = \Phi_3^*/6$ – symbols as in Figure 7.

implying the major evolution is a decline in space density (and global star formation rate) by $\simeq \times 6$ at $z \approx 6$. This sharp decline, which must represent a lower limit to the true decline given the likelihood of contamination from foreground sources, suggests it may be difficult for luminous star-forming i' -dropouts to be the main source of ionizing photons of the Universe.

We attempt to quantify this by comparing with the estimate of Madau, Haardt & Rees (1999) for the density of star formation required for reionization:

$$\dot{\rho}_{\text{SFR}} \approx 0.013 f_{\text{esc}}^{-1} \left(\frac{1+z}{6} \right)^3 \left(\frac{\Omega_b h_{50}^2}{0.08} \right)^2 C_{30} M_{\odot} \text{yr}^{-1} \text{Mpc}^{-3} \quad (2)$$

The escape fraction of ionizing photons (f_{esc}) for high-redshift galaxies is highly uncertain (e.g., Steidel, Pettini & Adelberger 2001), but even if we take $f_{\text{esc}} = 1$ (no absorption




Figure 9. The spatial distribution of our UDF i' -drops on the sky (diamonds). The location of the confirmed $z = 5.8$ source from Paper I is marked (#1) as are two other sources just outside the UDF, spectroscopically identified at $z = 5.8 - 5.9$ by Stanway et al. (2004a).

by H I) this estimate of the star formation density required is a factor of 3 higher than our measured star formation density of $0.005 h_{70}^{-2} M_{\odot} \text{yr}^{-1} \text{Mpc}^{-3}$ at $z \approx 6$ from galaxies in the UDF with SFRs $> 1 h_{70}^{-2} M_{\odot} \text{yr}^{-1}$. Unless we invoke another low-luminosity population (e.g., forming globular clusters; Ricotti 2002) or an extremely steep faint end slope ($\alpha \simeq -2$) then it would appear that these is insufficient UV flux from starburst at $z \approx 6$ to reionize the Universe. As AGN are also under-abundant at these epochs (e.g., Dijstra, Haiman & Loeb 2004) this presents challenge to the current thinking on the reionization of the Universe.

4 CONCLUSIONS

We summarize our main conclusions as follows:

(i) We present an i' -dropout catalogue of $z \approx 6$ star forming galaxy candidates in the Ultra Deep Field (UDF) to a limiting flux (8σ) of $z'_{AB} < 28.5$. This represents a substantial advance over the depths achieved in the GOODS catalogues and enables us, for the first time, to address questions concerning the likely form of the faint end of the luminosity function.

(ii) We detect 54 resolved sources with $(i' - z')_{AB} > 1.3$ in the deepest 11 arcmin² portion of the UDF and consider this to be an upper limit to the abundance of star forming galaxies at $z \approx 6$.

(iii) Exploiting the unique depth of the UDF, within the uncertainties of the small field we deduce that there is no evidence for a significant change in the form of the star forming luminosity function over $3 < z < 6$ other than in its absolute normalization.




Figure 10. An updated version of the ‘Madau-Lilly’ diagram (Madau et al. 1996; Lilly et al. 1996) illustrating the evolution of the comoving volume-averaged star formation rate. Our work from the UDF data is plotted a solid symbol. Other determinations have been recalculated for our cosmology and limiting UV luminosity of $\approx 1 h_{70}^{-2} M_{\odot} \text{ yr}^{-1}$ at $z \approx 6$ (equivalent to $\approx 0.1 L_{3}^*$ at $z \approx 3$ from Steidel et al. 1999), assuming a slope of $\alpha = -1.6$ for $z > 2$ and $\alpha = -1.3$ for $z < 2$. Data from the CFRS survey of Lilly et al. (1996) are shown as open circles; data from Connolly et al. (1997) are squares; and the Lyman break galaxy work of Steidel et al. (1999) is plotted as crosses, of Fontana et al. (2002) as inverted triangles and that by Iwata et al. (2003) as an open diamond. The upright triangles are the GOODS i' -drop results from Giavalisco et al. (2004). The three ACS estimates of Bouwens et al. (2003) are shown by small crossed circles and indicate three different completeness corrections for one sample of objects – the larger symbol is the recent re-determination using a new catalogue by this group from a deeper dataset (the UDF flanking fields – Bouwens et al. 2004); we have recomputed the comoving number density from the Bouwens et al. (2004) because of a discrepancy on the scale of their plot of star formation history (their Fig. 4 in astro-ph/0403167 v1 & v2).

(iv) Using simulations based on lower redshift data, we deduce that, regardless of contamination by foreground interlopers, the abundance of i' dropouts detected is significantly less than predicted on the basis of no evolution in the comoving star formation rate from $z = 3$ to $z = 6$. The UDF data supports our previous suggestions that the star formation rate at $z \approx 6$ was about $\times 6$ less than at $z \approx 3$ (Stanway, Bunker & McMahon 2003).

(v) The inferred comoving star formation rate of $0.005 h_{70}^2 M_{\odot} \text{ yr}^{-1} \text{ Mpc}^{-3}$ at $z \approx 6$ may pose a significant challenge for models which require that luminous star forming galaxies in the redshift range $6 < z < 10$ are responsible for reionizing the Universe.

(vi) The contamination of our i' -drop sample of candidate $z \approx 6$ galaxies by cool Galactic stars appears to be minimal at $z'_{AB} > 26$, possibly because we are seeing beyond the Galactic disk at the faint magnitudes probed by the UDF.

Acknowledgements

We thank colleagues at the Space Telescope Science Institute for making the UDF data available as a public database, on

schedule and in a manner suitable for immediate analysis. ERS acknowledges a Particle Physics and Astronomy Research Council (PPARC) studentship supporting this study.

REFERENCES

- Adelberger K. L., Steidel C. C., 2000, ApJ, 544, 218
 Barger A. J. et al., 2003, AJ, 126, 632
 Becker R. H. et al., 2001, AJ, 122, 2850
 Beckwith, S., Somerville, R., Stiavelli, M. 2003, STScI Newsletter vol 20 issue 04
 Bertin E., Arnouts S., 1996, A&AS, 117, 393
 Blanton M. R., et al., 2003, ApJ, 592, 819
 Bouwens R., et al., 2003, ApJ, 595, 589
 Bouwens R., et al., 2004, preprint (astro-ph/0403167)
 Bremer M. N., Lehnert M. D., Waddington I., Hardcastle M. J., Boyce P. J., Phillipps S., 2004, MNRAS, 347, L7
 Bunker A. J., Stanway E. R., Ellis R. S., McMahon R. G., McCarthy P. J., 2003, MNRAS, 342, L47 [Paper II]
 Connolly A. J., Szalay A. S., Dickinson M., Subbarao M. U., Brunner R. J., 1997, ApJ, 486, L11
 Cimatti A., et al., 2002, A&A, 381, L68
 Coleman G. D., Wu C.-C., Weedman D. W., 1980, ApJS, 43, 393
 Dickinson M., et al., 2004, ApJ, 600, L99
 Dijkstra M., Haiman Z., Loeb A., 2004, preprint (astro-ph/0403078)
 Efsthathiou G., Ellis R. S., Peterson B. A., 1988, MNRAS, 232, 431
 Fan X., et al., 2001, AJ, 122, 2833
 Fan X., et al., 2002, AJ, 123, 1247
 Ferguson H. C., et al., 2004, ApJ 600 L107
 Fontana A., Poli F., Menci N., Nonino M., Giallongo E., Cristiani S., D’Odorico S., 2003, ApJ, 587, 544
 Ford H. C., et al., 2002, BAAS, 200.2401
 Fruchter A., Hook R., 2002, PASP, 114, 144
 Gabasch A., et al., 2004, A&A in press, preprint (astro-ph/0403535)
 Giavalisco M., Dickinson M., 2002, ApJ, 550, 177
 Giavalisco M., et al., 2004, ApJ, 600, L103
 Gunn J. E., Peterson B. A., 1965, ApJ, 142, 1633
 Iwata I., Ohta K., Tamura N., Ando M., Wada S., Watanabe C., Akiyama M., Aoki K., 2003, PASJ, 55, 415
 Kogut A., et al., 2003, ApJS, 148, 161
 Lanzetta K. M., Yahata N., Pascarella S., Chen H.-W., Fernández-Soto A., 2002, ApJ, 570, 492
 Leitherer C., Heckman T. M., 1995, ApJS, 96, 9
 Lilly S. J., Tresse L., Hammer F., Crampton D., Le Fèvre O., 1995, ApJ, 455, 108
 Kennicutt R. C., 1998, ARA&A, 36, 189
 Koekemoer A. M., Fruchter A. S., Hook R. N., Hack W., 2002, HST Calibration Workshop, Eds. Arribas S., Koekemoer A. M., Whitmore B., (STScI: Baltimore), p. 325
 Madau P., Ferguson H. C., Dickinson M. E., Giavalisco M., Steidel C. C., Fruchter A., 1996, MNRAS, 283, 1388
 Madau P., Pozzetti L., Dickinson M., 1998, ApJ, 498, 106
 Madau P., Haardt F., Rees M., 1999, ApJ, 514, 648
 Meurer G. R., Heckman T. M., Lehnert M. D., Leitherer C., Lowenthal J., 1997, AJ, 114, 54
 Oke J. B., Korycansky D. G., 1982, ApJ, 225, 11
 Oke J. B., Gunn J. E., 1983, ApJ, 266, 713
 Ricotti M., 2002, MNRAS, 336, L33
 Salpeter E. E., 1955, ApJ, 121, 161
 Scalo J. M., 1986, Fund. Cosmic Phys., 11, 1
 Schlegel D. J., Finkbeiner D. P., Davis M., 1998, ApJ, 500, 525
 Somerville R. S., Lee K., Ferguson H. C., Gardener J. P., Moustakas L. A., Giavalisco M., 2004, ApJ, 600, L171

- Stanway E. R., Bunker A. J., McMahon R. G., 2003, MNRAS, 342, 439 [Paper I]
- Stanway E. R., et al., 2004a, ApJ, 604, L13
- Stanway E. R., Bunker A. J., McMahon R. G., Ellis R. S., Treu T., McCarthy P. J., 2004b, ApJ in press, preprint (astro-ph/0308124) [Paper III]
- Stanway, E. R., 2004, PhD Thesis (University of Cambridge)
- Stanway E. R., McMahon R. G., Bunker A. J., 2004c, submitted to MNRAS, preprint (astro-ph/0403585)
- Steidel C. C., Pettini M., Hamilton D., 1995, AJ, 110, 2519
- Steidel C. C., Giavalisco M., Pettini M., Dickinson M. E., Adelberger K. L., 1996, ApJ, 462, L17
- Steidel C. C., Adelberger K. L., Giavalisco M., Dickinson M. E., Pettini M., 1999, ApJ, 519, 1
- Steidel C. C., Pettini M., Adelberger K. L., 2001, ApJ, 546, 665
- Williams R. E., et al., 1996, AJ, 112, 1335
- Williams R. E., et al., 1998, BAAS, 30, 1366 (AAS 193, #75.01)
- Yan H., Windhorst R. A., Cohen S., 2003, ApJ, 585, L93
- Yan H., Windhorst R. A., 2004, ApJ, 600, L1

This paper has been typeset from a \TeX / \LaTeX file prepared by the author.